

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/10975950>

# Bioavailability of minerals in legumes

Article in *British Journal Of Nutrition* · January 2003

DOI: 10.1079/BJN/2002718 · Source: PubMed

---

CITATIONS

255

READS

3,141

1 author:



[Ann-Sofie Sandberg](#)

Chalmers University of Technology

210 PUBLICATIONS 6,874 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Herring and CVD risk factors [View project](#)



AIR-study [View project](#)

## Bioavailability of minerals in legumes

Ann-Sofie Sandberg\*

Department of Food Science, Chalmers University of Technology, PO Box 5401, SE 402 29 Göteborg, Sweden

The mineral content of legumes is generally high, but the bioavailability is poor due to the presence of phytate, which is a main inhibitor of Fe and Zn absorption. Some legumes also contain considerable amounts of Fe-binding polyphenols inhibiting Fe absorption. Furthermore, soya protein *per se* has an inhibiting effect on Fe absorption. Efficient removal of phytate, and probably also polyphenols, can be obtained by enzymatic degradation during food processing, either by increasing the activity of the naturally occurring plant phytases and polyphenol degrading enzymes, or by addition of enzyme preparations. Biological food processing techniques that increase the activity of the native enzymes are soaking, germination, hydrothermal treatment and fermentation. Food processing can be optimized towards highest phytate degradation provided that the optimal conditions for phytase activity in the plant is known. In contrast to cereals, some legumes have highest phytate degradation at neutral or alkaline pH. Addition of microbial enzyme preparations seems to be the most efficient for complete degradation during processing. Fe and Zn absorption have been shown to be low from legume-based diets. It has also been demonstrated that nutritional Fe deficiency reaches its greatest prevalence in populations subsisting on cereal- and legume-based diets. However, in a balanced diet containing animal protein a high intake of legumes is not considered a risk in terms of mineral supply. Furthermore, once phytate, and in certain legumes polyphenols, is degraded, legumes would become good sources of Fe and Zn as the content of these minerals is high.

### **Pulses: Minerals: Bioavailability: Phytate: Polyphenols: Oxalate**

Legumes are rich sources of food proteins from plants and have provided a protein source for man since the earliest civilizations. Legumes are often advocated in Western diets because of their beneficial nutritional effects and because they are a low-cost source of protein. Pulses represent dry grain legumes for human consumption. The content of Fe and other minerals is generally high in legumes (Table 1). However, the legumes also contain antinutritional factors, such as proteinase inhibitors, lectin, raffinose oligosaccharides, saponins, polyphenols and phytate. Antinutritional factors lower the nutritional value of a food by lowering the digestibility or bioavailability of nutrients. Phytate, and some of the degradation products of phytate, are well-known inhibitors of absorption of essential dietary minerals, in particular non-haem iron and Zn. Certain Fe-binding polyphenols are potent inhibitors of non-haem iron absorption.

The negative influence on Fe absorption is nutritionally the most important, especially in industrial products such as infant formulas, but more importantly in many developing countries where diet is based on cereal and legume products. Deficiency of Fe, and perhaps Zn, is highly prevalent in developing countries, but also in vulnerable groups with

high requirements in industrialized countries, such as women of fertile age, infants and adolescents. The increased number of vegetarians among young people might lead to increased prevalence of Fe deficiency, because the mineral availability may be crucial in a vegetarian diet (McEndree *et al.* 1983; Helman & Darnton-Hill, 1987; Reddy & Sanders, 1990). In developing countries Fe deficiency, due to poor bioavailability, retards normal brain development in infants and affects the success of a pregnancy by increasing premature deliveries, as well as morbidity of mother and child at or around childbirth. Zn deficiency prevents normal child growth and greatly weakens the immune system, leading to more infections. Nutritional Fe deficiency reaches its greatest prevalence and severity in populations subsisting predominantly on cereal and legume diets (International Nutritional Anemia Consultative Group, 1982).

### **Phytate**

Phytate (inositol hexaphosphate) constitutes 1–3% of cereal grains, legume seeds and nuts, and also occurs in low concentrations in roots, tubers and vegetables. In particular, wholegrain cereals and legumes have a high

---

\* Corresponding author: Dr Ann-Sofie Sandberg, tel +46 31 33 55 630, fax +46 31 83 37 82, email ann-sofie.sandberg@fsc.chalmers.se

**Table 1.** Minerals (iron, zinc, calcium and magnesium; per 100 g of dry seed) in some pulses and soyabeans (from Fachmann *et al.* 2000)

	Fe (mg)	Zn (mg)	Ca (mg)	Mg (mg)
<i>Phaseolus vulgaris</i>	7.0	3.0	197	250
Peas ( <i>Pisum sativum</i> )	7.36	3.01	96	132
Chickpeas	6.96	3.54	124	155
Lentils	7.50	3.73	71	129
Soyabeans	6.64	4.18	201	220

content of phytate but also of the minerals Zn, Fe and Mg. The phytate content of some legumes is shown in Table 2 and the mineral content in Table 1. In legume seeds, phytate is located in the protein bodies in the endosperm. Phytate occurs as a mineral complex, which is insoluble at the physiological pH of the intestine. It is considered anti-nutritional, causing reduced uptake in the human intestine of essential dietary minerals such as Fe, Zn and Ca. A dose-dependent inhibition of Fe, Zn and Ca absorption by phytate has been demonstrated in humans (Hallberg *et al.* 1989; Brune *et al.* 1992; Hurrell *et al.* 1992; Fredlund *et al.* 2002). Inositol pentaphosphate has also been identified as an inhibitor of Fe and Zn absorption (Lönnerdal *et al.* 1989; Sandström & Sandberg, 1992; Sandberg *et al.* 1999). Furthermore, it was found recently that inositol tri- and tetraphosphate contribute to the negative effect on Fe absorption of processed food containing a mixture of inositol phosphates (Sandberg *et al.* 1999), probably by interactions with the higher phosphorylated inositol phosphates.

### Polyphenols

Certain polyphenols are able to complex-bind Fe, which make the complex-bound Fe unavailable for absorption (Brune *et al.* 1989; Hurrell *et al.* 1999). According to Brune *et al.* (1989) the amount of Fe-binding phenolic gall-oil groups in foods roughly corresponds to the degree of inhibition of Fe absorption. Hurrell *et al.* (1999), however, concluded that all major types of food polyphenols can strongly inhibit dietary non-haem iron absorption.

Legumes contain varying amounts of polyphenols and generally the amounts are considered higher in the coloured seeds (reviewed by Salunkhe *et al.* 1982). Beans of the species *Phaseolus vulgaris* were found to contain high amounts of polyphenols (Paredes-Lopez & Harry, 1989), whereas the content of polyphenols in peas (*Pisum sativum*) is very low.

**Table 2.** Content of phytate-phosphorus in legumes (from Reddy *et al.* 1989)

Legumes	Phytate-P (g/100 g)
Soyabeans	0.28–0.63
Red kidney beans	0.34–0.58
Peas	0.06–0.33
Lentils	0.08–0.30

### Oxalic acid

Oxalate salts are poorly soluble at intestinal pH and oxalic acid is known to decrease Ca absorption in monogastric animals (Allen, 1982). The effect of oxalate on Ca absorption in humans is less clear. Ca in spinach with a high oxalate content is very poorly absorbed (Heaney *et al.* 1988) while kale, a low-oxalate vegetable, exhibits excellent Ca absorbability (Heaney & Weaver, 1990). However, studies with calcium oxalate suggest that other factors in spinach contribute to the low Ca absorption (Heaney & Weaver, 1989). A certain amount of oxalic acid also occurs in phaseolus beans.

### Enzymatic degradation of phytate in legumes during food processing

Degradation of phytate can occur both during food processing and in the gastrointestinal tract. This degradation is of nutritional importance because it has been demonstrated that such controlled degradation improves the uptake of essential dietary minerals, i.e. Fe and Zn (Sandberg & Svanberg, 1991; Brune *et al.* 1992; Hurrell *et al.* 1992). Major efforts are therefore made to reduce the amount of phytate in foods by means of phytate-degrading enzymes, phytases, present naturally in the plant foods or present in yeasts or other micro-organisms used in food processing. These enzymes successively remove, one after the other, the six phosphorus groups attached to the inositol ring. Biotechnologically produced microbial phytase preparations are now commercially available and used for feed preparations. In the future, their use in food processing could be feasible. In order to substantially increase Fe absorption, phytate degradation has to be virtually complete. Recent findings suggest that the inositol penta-, tetra- and triphosphates must also be degraded in order to improve Fe absorption (Sandberg *et al.* 1999).

However, ascorbic acid is a potent enhancer of Fe absorption, which can counteract the inhibitory effect of phytate (Hallberg *et al.* 1989; Siegenberg *et al.* 1991). Increasing the ascorbic acid in phytate-containing meals is thus another means of improving Fe absorption. Certain other organic acids formed during fermentation may also improve Fe absorption. Heat treatment of plant foods, on the other hand, often reduces the bioavailability due to inactivation of the enzyme phytase and destruction of ascorbic acid (reviewed by Sandberg, 1996).

Biological food-processing techniques that increase the activity of native enzymes of cereals and legumes are: soaking, germination, hydrothermal processing and fermentation. During germination, phytase enzymes are synthesized or activated. Lactic fermentation leads to lowering of pH as a consequence of bacterial production of organic acids, mainly lactic acid, which is favourable for cereal phytase activity (Sandberg, 2002). The micro-organisms (e.g. fungi) of the starter culture used in fermentation, in some cases, exert phytase activity. However, in contrast to fungi and yeast, *Lactobacillus* sp. was not found to produce phytase (Fredrikson *et al.* 2002b).

To optimize the food process to increase mineral availability by phytate degradation, it is essential to know the

optimal conditions for the phytases, which are responsible for phytate degradation in the food process. There are differences in optimal conditions for phytate degradation between plant species. Most cereal phytases have pH optima between 4.5 and 5.6, but pH optima of some legumes are neutral or alkaline (Loewus *et al.* 1990). Scott (1991) demonstrated an alkaline phytase extracted from different varieties of *Phaseolus vulgaris*. Gustafsson & Sandberg (1995) found that phytate degradation occurs in brown beans at pH 4.5 and at pH 8 at 37°C, but the highest phytate degradation was found at pH 7.0 and 55°C. By fermentation of pre-soaked beans, an 88% reduction of the phytate content was obtained.

In germinated or soaked lentils, faba beans and peas, the highest phytate degradation was found at 45°C and pH 5.0, 4.0 and 7.0, respectively (Honke *et al.* 1999). Fredrikson *et al.* (2001a) achieved the most efficient phytate degradation of pea flour at pH 7.5 and 45°C. At these soaking conditions it was possible to obtain an almost complete phytate degradation in 10 h, combined with 66% reduction of the sum of inositol penta-, tetra- and triphosphates. Favourable conditions for phytate degradation in black beans were found at pH 5.5 and 50°C (Greiner & Konietzny, 1997). However, compared to soaking at the mentioned conditions, a more extensive degradation was obtained by adding exogenous phytase.

#### Enzymic degradation of polyphenols in legumes during processing

Enzymic degradation of polyphenols during processing should also be a possible strategy for improvement of Fe availability. Moreover, the addition of ascorbic acid was found to prevent the dose-dependent inhibitory effect of polyphenols on non-haem iron absorption (Siegenberg *et al.* 1991).

At least for high-tannin cereals, it has been shown that treatment with polyphenol oxidase had a reducing effect on the phenolic content (Matuschek *et al.* 2001). Addition of 1500 U/g resulted in a 60% reduction of phenolic compounds and a significant improvement of Fe availability, estimated *in vitro*. A fungal tannase was used to decompose phenolic compounds in brown beans, but the influence on Fe availability was not determined (Gustafsson & Sandberg, 1995). Germination and fermentation of lentils were found to modify the phenolic composition. Germination reduced the amount of phenolic compounds, while natural fermentation was found to increase certain phenolic monomers (catechin) (Bartolomé *et al.* 1997).

#### Iron and zinc absorption from legume-based diets

Fe absorption from soyabeans and soya protein products studied in single meals, using extrinsic labelling of meals with radioactive isotopes or studied by a stable isotope technique, was found to be low (Cook *et al.* 1981; Morck *et al.* 1981; Hallberg & Rossander, 1982; Hurrell *et al.* 1992; Davidsson *et al.* 1994). Fe absorption from single meals based on black beans, lentils, mung beans and split beans was found to be very low, ranging from 0.8% to 1.9% (Lynch *et al.* 1984).

Studies of meals based on white beans and soyabeans indicate that the effect of phytate on Zn absorption is less pronounced in soyabeans than in cereals (Sandström & Cederblad, 1980; Sandström *et al.* 1989) and that legume and animal protein are comparable, with the same Zn content. Low Zn absorption from a soya-protein-based infant formula has, however, been found (Sandström *et al.* 1983).

The enzymatic degradation of phytate, by addition of a microbial phytase preparation, in soya infant formula was found to improve Fe absorption significantly, provided that the removal of phytate was virtually complete (Hurrell *et al.* 1992). However, in the same study it was found that even after removal of phytate, soya protein itself is still relatively inhibitory to Fe absorption, probably due to the presence of Fe-binding peptides. Davidsson *et al.* (1994) demonstrated that the Fe bioavailability can be increased by either removal of phytate or increasing the ascorbic acid content in soya-based infant formulas. It has also been demonstrated that removal of phytate from soya formulas significantly improves Zn absorption (Lönnerdal *et al.* 1988).

#### The New Technologies for Improved Nutritional and Functional Value of Pea Protein project

The general objective of the EU-funded project, NUT-RIPEA (FAIR CT 95-0193), was to use new technologies to develop improved pea protein products, which are devoid of antiphenological and antinutritional factors. The nutritional and functional properties of pea proteins suggest a high potential for use in food products. Therefore, the purpose of the project was to design and develop a technical process: (1) to prepare improved pea protein products under pilot plant and factory conditions; (2) to evaluate the functional and sensory properties of improved pea products added to a variety of foods for human consumption; (3) to screen *in vitro* and in animal models the nutritional properties and antigenicity of the protein products; (4) to develop an infant formula based on the improved pea protein products; and (5) to evaluate the formula for antigenicity and protein quality in animals and Fe absorption in humans. The results have been published recently (Sandberg, 2000).

The use of pea protein isolate could be an alternative to soya isolate. Soya formulas have been used for a long period and the nutritional status of infants fed soya formula has been well documented and found to be similar to that of infants fed cow's milk formulas. However, the bioavailability of nutrients, especially minerals, has been reported to be lower than that of milk-based formulas. The availability of Fe and Zn in a dephytinized infant formula based on pea protein was evaluated (Fredrikson *et al.* 2001b, 2002a). Soluble amounts of Fe and Zn were collected during simulated *in vitro* digestion performed in a computer-controlled dynamic gastrointestinal model. The results showed that the complete dephytinization of pea protein, by addition of a microbial phytase preparation during processing, increased the amount of Fe and Zn potentially available for absorption, by 50% and 100%, respectively. Also, an improved uptake of Fe

was demonstrated *in vitro* in Caco 2 cells (Fredrikson *et al.* 2002a). The improvement of Fe availability was confirmed in a human study, the Fe absorption increasing by more than 50% after phytate removal or addition of ascorbic acid to the pea protein infant formula (Davidsson *et al.* 2001). The data in the human study also suggest that fractional Fe absorption is higher from the dephytinized pea protein formula compared to dephytinized soya protein formulas. Pea protein may therefore be a nutritional beneficial alternative to soya protein in infant formulas, but further evaluation is needed.

### Conclusions

The absorption of minerals depends on the total composition of the meal. In a balanced diet containing animal protein, a high intake of legumes does not imply a risk of inadequate mineral supply. Strictly vegetarian diets based on unrefined cereals and possibly also some legume-based diets will result in low absorption of Fe and Zn. However, the utilization of Fe and Zn, and probably other minerals, can be improved by food processes such as fermentation with phytase-producing microorganisms; soaking and germination, which degrade phytate; or addition of phytase and, in some legumes, polyphenol-degrading enzymes. In the modern food industry, the phytate content of soya-based infant formulas is of concern; major efforts are therefore being made to remove phytate. Once phytate is degraded, legumes would become good sources of Fe and Zn as the content of these minerals is high.

### References

- Allen LK (1982) Calcium bioavailability and absorption; a review. *American Journal of Clinical Nutrition* **35**, 783–808.
- Bartolomé B, Hernández T & Estrella I (1997) Effects of processing on individual condensed tannins from lentils. In *COST 98 Effects of Antinutrients on the Nutritional Value of Legume Diets*, vol. 4, pp. 32–36 [S Bardocz, M Muzquiz and A Pusztai, editors]. Luxembourg: European Communities.
- Brune M, Rossander L & Hallberg L (1989) Iron absorption and phenolic compounds. Importance of different phenolic structures. *European Journal of Clinical Nutrition* **43**, 547–558.
- Brune M, Rossander-Hulthén L, Hallberg L, Gleerup A & Sandberg A-S (1992) Human iron absorption from bread: Inhibiting effects of cereal fiber, phytate and inositol phosphates with different numbers of phosphate groups. *Journal of Nutrition* **122**, 442–449.
- Cook JD, Morck TA & Lynch SR (1981) The inhibitory effect of soy products on nonheme iron absorption in man. *American Journal of Clinical Nutrition* **34**, 2622–2629.
- Davidsson L, Dimitriou T, Walczyk T & Hurrell RF (2001) Iron absorption from experimental infant formulas based on pea protein isolate. The effect of phytic acid and ascorbic acid. *British Journal of Nutrition* **85**, 59–63.
- Davidsson L, Galan P, Kastenmayer P, Cherouvrier F, Juillerat M-A, Jerberg S & Hurrell RF (1994) Iron bioavailability studied in infants: The influence of phytic acid and ascorbic acid in infant formulas based on soy isolate. *Pediatric Research* **36**, 816–822.
- Fachmann W, Souci SW & Kraut H (2000) *Food Composition and Nutrition Tables*, p. 1182. Boca Raton: CRC Press.
- Fredlund K, Rossander-Hulthén L, Isaksson M, Almgren A & Sandberg AS (2002) Absorption of zinc and calcium: dose-dependent inhibition by phytate. *Journal of Applied Microbiology* **93**, 197–204.
- Fredrikson M, Alminger ML, Carlsson NG & Sandberg A-S (2001a) Phytate content and phytate degradation by endogenous phytase in pea (*Pisum sativum*). *Journal of the Science of Food and Agriculture* **81**, 1139–1144.
- Fredrikson M, Alminger ML & Sandberg AS (2002a) Improved *in vitro* availability of iron and zinc from dephytinized pea protein formulas, comparison of iron availability with commercial soy protein formula. Submitted for publication.
- Fredrikson M, Andlid T, Haikara A & Sandberg AS (2002b) Phytate degradation by microorganisms in synthetic media and pea flour. *Journal of Applied Microbiology* **93**, 197–204.
- Fredrikson M, Biot P, Alminger ML, Carlsson NG & Sandberg AS (2001b) Production process for high-quality pea-protein isolate, with low content of oligosaccharides and phytate. *Journal of Agricultural and Food Chemistry* **49**, 1208–1212.
- Greiner R & Konietzny U (1997) Phytate hydrolysis in black beans by endogeneous and exogeneous enzymes. In *COST 98 Effects of Antinutrients on the Nutritional Value of Legume Diets*, vol. 4, pp. 19–27 [S Bardocz, M Muzquiz and A Pusztai, editors]. Luxembourg: European Communities.
- Gustafsson E & Sandberg A-S (1995) Phytate reduction in brown beans (*Phaseolus vulgaris* L.). *Journal of Food Science* **60**, 149–152, 156.
- Hallberg L, Brune M & Rossander L (1989) Iron absorption in man: ascorbic acid and dose-dependent inhibition by phytate. *American Journal of Clinical Nutrition* **49**, 140–144.
- Hallberg L & Rossander L (1982) Effect of soy protein on non-heme iron absorption in man. *American Journal of Clinical Nutrition* **36**, 514–520.
- Heaney RP & Weaver CM (1989) Oxalate: effect on calcium absorbability. *American Journal of Clinical Nutrition* **50**, 830–832.
- Heaney RP & Weaver CM (1990) Calcium absorption from kale. *American Journal of Clinical Nutrition* **51**, 656–657.
- Heaney RP, Weaver CM & Recker RR (1988) Calcium absorption from spinach. *American Journal of Clinical Nutrition* **47**, 707–709.
- Helman AD & Darnton-Hill I (1987) Vitamin and iron status in new vegetarians. *American Journal of Clinical Nutrition* **45**, 785–789.
- Honke J, Sandberg A-S & Kozłowska H (1999) The influence of pH and temperature on endogenous phytase activity and on hydrolysis of inositol hexaphosphate in lentil, faba bean and pea seeds. *Polish Journal of Food and Nutrition Sciences* **8/49**, 109–122.
- Hurrell RF, Juillerat M-A, Reddy MB, Lynch SR, Dassenko SA & Cook JD (1992) Soy protein, phytate, and iron absorption in humans. *American Journal of Clinical Nutrition* **56**, 573–578.
- Hurrell RF, Reddy M & Cook JD (1999) Inhibition of non-haem iron absorption in man by polyphenolic-containing beverages. *British Journal of Nutrition* **81**, 289–295.
- International Nutritional Anemia Consultative Group (1982) *Iron Absorption from Cereals and Legumes. A Report of the International Nutritional Anemia Consultative Group*, New York, pp. 1–44. New York: The Nutrition Foundation.
- Loewus RA, Everard JD & Young KA (1990) 3. Inositol metabolism: Precursor role and breakdown. In *Inositol Metabolism in Plants*, pp. 21–45 [DJ Morre, WF Boss and FA Loewus, editors]. New York: Wiley-Liss.
- Lönnerdal B, Bell JG, Hendricks AG, Burns RA & Keen CL

- (1988) Effect of phytate removal on zinc absorption from soy formula. *American Journal of Clinical Nutrition* **48**, 1301–1306.
- Lönnerdal B, Sandberg A-S, Sandström B & Kunz C (1989) Inhibitory effects of phytic acid and other inositol phosphates on zinc and calcium absorption in suckling rats. *Journal of Nutrition* **119**, 211–214.
- Lynch SR, Beard JL, Dassenko SA & Cook JD (1984) Iron absorption from legumes in humans. *American Journal of Clinical Nutrition* **40**, 42–47.
- McEndree LS, Kies CV & Fox HM (1983) Iron intake and iron nutritional status of lacto-ovo-vegetarian and omnivore students eating in a lacto-ovo-vegetarian food service. *Nutrition Report International* **27**, 199–206.
- Matuschek E, Towo E & Svanberg U (2001) Oxidation of polyphenols in high-tannin cereals and the effect on iron bioavailability. In *Bioavailability 2001. Abstract book*, [B Abt, R Amadò and L Davidsson, editors]. Zürich: ETH Swiss Federal Institute of Technology.
- Morck TA, Lynch SR, Skikne BS & Cook JD (1981) Iron availability from infant food supplements. *American Journal of Clinical Nutrition* **34**, 2630–2634.
- Paredes-Lopez O & Harry GI (1989) Changes in selected chemical and antinutritional components during tempeh preparation using fresh and hardened common beans. *Journal of Food Science* **54**, 968–970.
- Reddy NR, Pierson MD, Sathé SK & Salunkhe DK (1989) Occurrence, distribution, content, and dietary intake of phytate. In *Phytates in Cereals and Legumes*, pp. 39–56 [NR Reddy, MD Pierson, SK Sathé and DK Salunkhe, editors]. Boca Raton, Florida: CRC Press.
- Reddy S & Sanders TAB (1990) Haematological studies on premenopausal Indian and Caucasian vegetarians compared with Caucasian omnivores. *British Journal of Nutrition* **64**, 331–338.
- Salunkhe DK, Jadhav SJ, Kadam SS & Chavan JK (1982) Chemical, biochemical, and biological significance of polyphenols in cereals and legumes. *CRC Critical Reviews in Food Science and Nutrition* **17**, 277–305.
- Sandberg AS (1996) Food processing influencing iron bioavailability. In *Iron Nutrition in Health and Disease*, pp. 349–356 [H Hallberg and N-G Asp, editors]. London: John Libbey.
- Sandberg AS (2000) Developing functional ingredients. A case study. In *Functional Foods*, pp. 209–232 [GR Gibson and CM Williams, editors]. Cambridge and Boca Raton, Florida: Woodhead Publishing and CRC Press.
- Sandberg AS (2002) *In vitro* and *in vivo* degradation of phytate. In *Food Phytates*, pp. 139–155 [NR Reddy and SK Sathé, editors]. Boca Raton, Florida: CRC Press.
- Sandberg AS, Brune M, Carlsson NG, Hallberg L, Skoglund E & Rossander-Hulthén L (1999) Inositol phosphates with different number of phosphate groups influence iron absorption in humans. *American Journal of Clinical Nutrition* **70**, 240–246.
- Sandberg AS & Svanberg U (1991) Phytate hydrolysis by phytase in cereals. Effects on *in vitro* estimation of iron availability. *Journal of Food Science* **56**, 1330–1333.
- Sandström B, Almgren A, Kivistö C & Cederblad A (1989) Effect of protein level and protein source on zinc absorption in man. *Journal of Nutrition* **119**, 48–53.
- Sandström B & Cederblad A (1980) Zinc absorption from composite meals. II. Influence of the main protein source. *American Journal of Clinical Nutrition* **33**, 1778–1783.
- Sandström B, Cederblad A & Lönnerdal B (1983) Zinc absorption from human milk, cow's milk and infant formulas. *American Journal of Diseases of Children* **137**, 726–729.
- Sandström B & Sandberg A-S (1992) Inhibitory effects of isolated inositol phosphates on zinc absorption in humans. *Journal of Trace Elements and Electrolytes in Health and Disease* **6**, 99–103.
- Scott JJ (1991) Alkaline phytase activity in nonionic detergent extracts of legume seeds. *Plant Physiology* **95**, 1298–1301.
- Siegenberg D, Baynes RD, Bothwell TH, Macfarlane BJ, Lamparelli RD, Car NG, MacPhail P, Schmidt U, Tal A & Mayet F (1991) Ascorbic acid prevents the dose-dependent inhibitory effects of polyphenols and phytates on nonheme-iron absorption. *American Journal of Clinical Nutrition* **53**, 537–541.